

FAST TRACK COMMUNICATION

Statistical correlations in a Coulomb gas with a test charge**Henning Schomerus**

Department of Physics, Lancaster University, Lancaster, LA1 4YB, UK

Received 16 June 2008, in final form 27 June 2008

Published 17 July 2008

Online at stacks.iop.org/JPhysA/41/332002**Abstract**

A recent paper (Jokela *et al* 2008 *Preprint* arXiv:0806.1491) contains a surmise about an expectation value in a Coulomb gas which interacts with an additional charge ξ that sits at a fixed position. Here, I demonstrate the validity of the surmised expression and extend it to a certain class of higher cumulants. The calculation is based on the analogy to statistical averages in the circular unitary ensemble of random-matrix theory and exploits properties of orthogonal polynomials on the unit circle.

PACS numbers: 02.50.Sk, 05.20.-y, 05.30.Fk, 05.45.Mt

1. Purpose and result

In a recent paper Jokela, Järvinen and Keski-Vakkuri studied n -point functions in timelike boundary Liouville theory via the analogy to a Coulomb gas on a unit circle [1]. In this analogy, N unit charges at position t_i interact with additional charges of integer value ξ_a , situated at position τ_a . To illustrate this technique the authors of [1] considered the canonical expectation value

$$\langle \cdot \rangle \equiv \frac{1}{Z} \int \prod_{i=1}^N \frac{dt_i}{2\pi} \prod_{i < j} |e^{it_i} - e^{it_j}|^2 \prod_i |e^{i\tau} - e^{it_i}|^{2\xi} (\cdot) \quad (1)$$

(where Z is a normalization factor so that $\langle 1 \rangle = 1$) and surmised that

$$\langle \text{Re } a_1 \rangle \equiv \left\langle \sum_i \cos(\tau - t_i) \right\rangle = -\frac{\xi N}{N + \xi}. \quad (2)$$

In this communication I demonstrate the validity of (2) and also compute expectation values of the more general quantities

$$a_n \equiv \sum_{i_1 < i_2 < \dots < i_n} \exp\left(i \sum_{k=1}^n (t_{i_k} - \tau)\right). \quad (3)$$

As a result, I find

$$\langle a_n \rangle = (-1)^n \frac{(N - n + 1)^{(\xi)} (n + 1)^{(\xi-1)}}{(N + 1)^{(\xi)} (1)^{(\xi-1)}} \quad \forall \quad n = 1, 2, \dots, N, \quad \xi \geq 0, \quad (4)$$

where $(x)^{(y)} = \Gamma(x + y) / \Gamma(x)$ is the generalized rising factorial (Pochhammer symbol). In particular, the validity of (2) follows from (4) by setting $n = 1$.

Expression (4) will be obtained by relating the generating polynomial

$$\varphi_{N,\xi}(\lambda) \equiv \sum_{n=0}^N \langle a_n \rangle (-\lambda)^{N-n} \quad (5)$$

to a weighted average of the secular polynomial in the circular unitary ensemble (CUE). This in turn establishes a relation to the Szegő polynomial of a Toeplitz matrix composed of binomial coefficients. This calculation sidesteps Jack polynomials and generalized Selberg integrals, which can be used to tackle general expectation values in multicomponent Coulomb gases [2].

2. Reformulation in terms of random matrices

The CUE is composed of $(N \times N)$ -dimensional unitary matrices U distributed according to the Haar measure. Identify t_i with the eigenphases of such a matrix. The joint probability distribution is then given by [3]

$$P(\{t_i\}_{i=1}^N) = z \prod_{i < j} |e^{it_i} - e^{it_j}|^2, \quad (6)$$

where z is again a normalization constant. This expression can also be written as the product of two Vandermonde determinants $\det V^+ \det V^-$ with matrices $V_{lm}^\sigma = e^{i\sigma(m-1)t_l}$. Furthermore, we can write

$$\prod_i |e^{i\tau} - e^{it_i}|^{2\xi} = [\det(1 - U e^{-i\tau}) \det(1 - U^\dagger e^{i\tau})]^\xi. \quad (7)$$

Finally, the expressions a_n in (3) arise as the expansion coefficients of the secular polynomial

$$\det(U e^{-i\tau} - \lambda) = \sum_{n=0}^N a_n (-\lambda)^{N-n}. \quad (8)$$

Note that in all these expressions τ can be shifted to any fixed value by a uniform shift of all t_i 's, which leaves the unitary ensemble invariant. Therefore, the expectation values are independent of τ . Collecting all results, we have the identity

$$\varphi_{N,\xi}(\lambda) = \frac{\langle [\det(1 - U) \det(1 - U^\dagger)]^\xi \det(U - \lambda) \rangle_{\text{CUE}}}{\langle [\det(1 - U) \det(1 - U^\dagger)]^\xi \rangle_{\text{CUE}}}. \quad (9)$$

This can be interpreted as a weighted average of the secular polynomial in the CUE.

3. Random-matrix average

Statistical properties of the secular polynomial without the weight factor ($\xi = 0$) have been considered in [4]. Clearly, $\varphi_{N,0} = (-\lambda)^N$, so that in this case the attention quickly moves on to higher moments of a_n . The main technical observation in [4] which allows us to address the case of finite ξ concerns averages of expressions $g(\{t_i\}_{i=1}^N)$ that are completely symmetric in all eigenphases. In this situation the average can be found via

$$\langle g(\{t_i\}_{i=1}^N) \rangle_{\text{CUE}} = \int \prod_i \frac{dt_i}{2\pi} g(\{t_i\}_{i=1}^N) \det W, \quad (10)$$

where $W_{lm} = e^{it_m(l-m)}$. Equation (10) is simpler than the general expression involving the product of two Vandermonde matrices, since each eigenphase only appears in a single column of W .

In the present problem, the numerator in (9) is represented by the completely symmetric function

$$g_1(\{t_i\}_{i=1}^N) = \prod_{i=1}^N [(e^{it_i} - \lambda)(1 - e^{it_i})^\xi(1 - e^{-it_i})^\xi], \tag{11}$$

while for the denominator we need to consider the similar expression

$$g_2(\{t_i\}_{i=1}^N) = \prod_{i=1}^N [(1 - e^{it_i})^\xi(1 - e^{-it_i})^\xi]. \tag{12}$$

Using the multilinearity of the determinant we can now pull each factor into the i th column and perform the integrals. This delivers the representation

$$\varphi_{N,\xi}(\lambda) = \frac{\det(B - \lambda A)}{\det A}, \tag{13}$$

where the matrices $A_{lm} = (-1)^{l-m} \binom{2\xi}{\xi+l-m}$, $B_{lm} = (-1)^{l-m+1} \binom{2\xi}{\xi+l-m+1}$ have entries given by binomial coefficients. We now exploit the regular structure of these matrices in two steps.

- (i) Matrix B contains the same entries as matrix A , but shifted to the left by one column index. In order to exploit this, let us expand the determinant in the numerator into a sum of determinants of matrices labeled by $X = (x_m)_{m=1}^N$, where we select each column either from A ($x_m = A$) or from B ($x_m = B$). (Note that we set these symbols in roman letters.) The related structure of A and B then entails that $\det X$ vanishes if X contains a subsequence $(x_m, x_{m+1}) = (A, B)$. Consequently we only need to consider determinants of matrices $X_n \equiv (B)_{m=1}^n \oplus (A)_{m=n+1}^N$, associated with sequences that contain n leading B's and $N-n$ trailing A's. As A is multiplied by $-\lambda$, $\det X_n$ contributes to order $(-\lambda)^{N-n}$. (Note that $X_0 = A$ and $X_N = B$.)
- (ii) Next, consider the matrix A_{N+1} , where the subscript denotes the dimension, and strike out the first row and the $n+1$ st column ($n = 0, 1, 2, \dots, N$). This takes exactly the form of the matrix X_n of dimension N . Therefore, the expressions $(-1)^n \det X_n$ are the cofactors of the first row of A_{N+1} . These, in turn, are proportional to the first column of A_{N+1}^{-1} , where the proportionality factor is given by $\det A_{N+1}$. Consequently, taking care of all alternating signs,

$$\varphi_{N,\xi}(\lambda) = (-1)^N \frac{\det A_{N+1}}{\det A_N} \sum_{n=0}^N (A_{N+1}^{-1})_{1,1+n} \lambda^{N-n}. \tag{14}$$

Via steps (i) and (ii) we have eliminated any reference to the matrix B .

4. Orthogonal polynomials

Matrix A is a Toeplitz matrix, $A_{lm} = c_{l-m}$. In order to find the explicit expression (4) we now make contact to the theory of orthogonal polynomials on the unit circle [5]. Among its many applications, this theory provides a general expression for the inverse of any Toeplitz matrix in terms of Szegő polynomials $\psi_N(\lambda)$. For the case of real symmetric coefficients, the inverse is generated via

$$\frac{\lambda \mu^N \psi_N(\lambda) \psi_N(\mu^{-1}) - \lambda^N \mu \psi_N(\lambda^{-1}) \psi_N(\mu)}{\lambda - \mu} = \frac{\det A_{N+1}}{\det A_N} \sum_{n,m=0}^N (A_{N+1}^{-1})_{m+1,n+1} \lambda^{N-n} \mu^m. \tag{15}$$

Comparison of this equation with $m = 0$ to (14) immediately leads to the identification of $(-1)^N \varphi_{N,\xi}(\lambda)$ with the Szegő polynomial $\psi_N(\lambda)$ of degree N . These polynomials satisfy recursion relations which for real symmetric coefficients take the form

$$\gamma_N = -\frac{1}{\delta_{N-1}} \oint \frac{d\lambda}{2\pi i} \psi_{N-1}(\lambda) \sum_{n=-\infty}^{\infty} c_n \lambda^n, \quad (16a)$$

$$\psi_N(\lambda) = \lambda \psi_{N-1}(\lambda) + \gamma_N \lambda^{N-1} \psi_{N-1}(\lambda^{-1}), \quad (16b)$$

$$\delta_N = \delta_{N-1} (1 - \gamma_N^2). \quad (16c)$$

The initial conditions are $\delta_0 = c_0$, $\psi_0(\lambda) = 1$. The numbers γ_N are known as the Schur or Verblunsky coefficients.

It can now be seen in an explicit if tedious calculation that the polynomials

$$\psi_N(\lambda) = (-1)^N \varphi_{N,\xi}(\lambda) = \sum_{n=0}^N \frac{(N-n+1)^{\binom{\xi}{n}} (n+1)^{\binom{\xi-1}{n}}}{(N+1)^{\binom{\xi}{n}} (1)^{\binom{\xi-1}{n}}} \lambda^{N-n} \quad (17a)$$

$$= \lambda^N {}_2F_1(-N, \xi; -N - \xi; \lambda^{-1}) \quad (17b)$$

(with coefficients and expansion given in (4) and (5)) indeed fulfil the Szegő recursion generated by the binomial coefficients $c_n = (-1)^n \binom{2\xi}{\xi-n}$. The recursion coefficients take the simple form

$$\gamma_N = \frac{\xi}{\xi + N}, \quad \delta_N = \frac{N!(2\xi + 1)^{\binom{N}{2}}}{[(\xi + 1)^{\binom{N}{2}}]^2}. \quad (17c)$$

This completes the proof of (4) and also entails the validity of (2).

Acknowledgments

The author wishes to thank Niko Jokela, Matti Järvinen and Esko Keski-Vakkuri for useful correspondence regarding [1], and Mitsuhiro Arikawa and Peter Forrester for useful communications about [2]. This work was supported by the European Commission, Marie Curie Excellence Grant MEXT-CT-2005-023778 (Nanoelectrophotonics).

References

- [1] Jokela N, Järvinen M and Keski-Vakkuri E 2008 *Preprint* [arXiv:0806.1491](https://arxiv.org/abs/0806.1491)
- [2] Forrester P J 1992 *J. Phys. A: Math. Gen.* **25** L607
Forrester P J 1992 *Nucl. Phys. B* **388** 671
- [3] Mehta M L 1991 *Random Matrices* 2nd edn (New York: Academic)
- [4] Haake F, Kuś M, Sommers H-J, Schomerus H and Życzkowski K 1996 *J. Phys. A: Math. Gen.* **29** 3641
- [5] Simon B 2005 *Orthogonal Polynomials on the Unit Circle: Part 1. Classical Theory; Part 2: Spectral Theory* (Providence, RI: American Mathematical Society)